Application No. 10/715,659 Docket No.: 2032(227045)

AMENDMENTS TO THE CLAIMS

1. - 7. (Cancelled)

8. (Currently amended) A method for treating an animal with a Th1 or Th2 related disease and monitoring the efficacy of said treatment comprising

administering a helminthic parasite preparation that alters a regulatory T cell activity to said animal, wherein said administering alters the activity of regulatory T cells; and

measuring said regulatory T cell activityresponses;

wherein an increase in the level of the regulatory T cell activity after said administering is indicative of the efficacy of said treatment.

9.-16. (Cancelled)

- 17. (Currently amended) The method of claim 8, wherein said regulatory T cell <u>activity</u> responses are is measured by determining the level of a regulatory T cell marker.
- 18. (Previously presented) The method of claim 17, wherein said regulatory T cell marker is an internal marker.
- 19. (Previously presented) The method of claim 18, wherein said internal marker is Scurfin, Smad7, Gata3, or Tbet (Tbx21).
- 20. (Previously presented) The method of claim 17, wherein said regulatory T marker is a cell surface marker.
- 21. (Previously presented) The method of claim 20, wherein said cell surface marker is selected from the group consisting of: CD4, CD45RB^{lo}, CD45Rc, Cytotoxic T lymphocyte associated antigen 4 (CTLA-4), Ox40, 4-1BB, CD25, CD103, CD62L, α_Eβ integrin, latency-

Application No. 10/715,659 Docket No.: 2032(227045)

associated peptide (LAP) or glucocorticoid induced TNF receptor family related protein (GITR), chemokine receptor CCR5, TI-ST2.

- (Previously presented) The method of claim 17, wherein said regulatory T cell marker is a secreted marker.
- (Previously presented) The method of claim 22, wherein said secreted marker is selected from the group consisting of IL-4, IL-13, IL-5, IL-10 or TGFβ, IFNγ and PgE2.
- 24. (Cancel IL-10)
- 25. (Currently amended) The method of claim 8.23_7 wherein said increase in the level of regulatory T cell activity is reflected by secretion by said regulatory T cell of secretes at least a 2-fold increase of TGF β as compared to naive T cells.
- 26. (Currently amended) The method of claim 8,23, wherein said increase in the level of regulatory T cell activity is reflected by secretion by said regulatory T cell of secretes at least a 2-fold less IFNy as compared to naive T cells.